

Comment on "Magnetic Breakdown at High Fields: Semiclassical and Quantum Treatments"

Keita Kishigi and Yasumasa Hasegawa

Faculty of Science, Himeji Institute of Technology, Ako, Hyogo 678-1297, Japan

(Received 18 September, 2000)

PACS numbers: 71.18.+y, 71.20.Rv, 72.15.Gd

In a recent Letter, Han *et al.* [1] investigated the effects of finite temperature and spin on the de Haas-van Alphen (dHvA) oscillation in the two-dimensional model, where the closed orbit and open orbit exist. They obtained the effective mass from the fitting of the g_e -dependence of these FTAs with the damping factor,

$$R_s = \cos(p\pi g_e m^*/2m_e), \quad (1)$$

expected from the semiclassical description based on the Lifshitz-Kosevich formula [2]. In this Comment we argue that since the damping factor Eq.(1) is obtained for the system with fixed chemical potential μ , it is not appropriate for the system with the fixed electron number N , although the estimation of m^* is the same. We also call attention that the spin does not reduce all FTAs but enhances some amplitudes in the system with magnetic breakdown. This interesting effect of spin [3] is seen in Fig. 6 of Ref. [1] but it was not mentioned.

For simplicity we first consider the two-dimensional free-electron system. In the case of fixed μ , the magnetization oscillates as a 'saw-tooth' given by [2],

$$\frac{M}{\beta N_0} = - \sum_{p=1}^{\infty} \frac{\sin\left(2\pi p \left(\frac{\mu}{\beta H} - \frac{1}{2}\right)\right)}{\pi p}, \quad (2)$$

where $\beta = e\hbar/m^*c$ is a double Bohr magneton with effective mass and N_0 is the total number of electrons at $H = 0$. When the spin-splitting is taken into account, the dHvA oscillation is just the sum of the contributions from spin-up and spin-down electrons with the phase shift of $\pm g_e m^*/4m_e$, as shown in Fig. 1(a), resulting in the damping factor Eq.(1). In the fixed N case, on the other hand, the magnetization for $g_e = 0$ is given by [2],

$$\frac{M}{\beta N} = \sum_{p=1}^{\infty} \frac{\sin\left(2\pi p \frac{N}{2\rho\beta H}\right)}{\pi p}, \quad (3)$$

where ρ is the density of states for each spin at $H = 0$. When the spin-splitting is considered, the chemical potential jumps between a spin-up Landau level and a spin-down Landau level when $N/2\rho\beta H$ is integer or integer plus 1/2. As a result, $\mp 2(g_e m^*/2m_e - [g_e m^*/2m_e])$, where $[g_e m^*/2m_e]$ is the integer part of $g_e m^*/2m_e$, is added to $M/\beta N$ when $n < N/2\rho\beta H < n + 1/2$ or $n + 1/2 < N/2\rho\beta H < n + 1$, respectively, as shown in Fig. 1(b). This modification of the magnetization is written by the damping factor as [3,4],

$$R_p^N = \begin{cases} |1 - 2(g_e m^*/2m_0 - [g_e m^*/2m_0])| & \text{for odd } p \\ 1 & \text{for even } p \end{cases} \cdot (4)$$

Only for the fundamental frequency ($p = 1$), the periods of the damping factor with respect to g_e for fixed μ (Eq. (1)) and fixed N (Eq.(4)) are the same. The g_e -dependence of the peak amplitude in Fig. 6 of Ref. [1] was fitted with the cosine curves (Eq.(1)), but it (especially for f_α) can be fitted better with Eq.(4).




FIG. 1. Magnetization as a function of the inverse magnetic field. (a) μ is fixed. (b) N is fixed. Thin and thick lines are the magnetization for $g_e = 0$ and $g_e \neq 0$, respectively.

The *enhancement* of the peak amplitude due to spin is seen for the $f_{\beta+\alpha}$ and $f_{2\alpha}$ in Fig.6 of Ref. [1]. The enhancement of $f_{\beta+\alpha}$ has been reported previously [3], where slightly simpler but essentially the same model is studied. We interpreted the enhancement of the amplitude as the result of incomplete cancellation between the effects of the chemical potential oscillation and magnetic breakdown [3]. We also observe the enhancement of the $\beta + 2\alpha$ oscillation [3]. The g_e -dependence of the 2α oscillation in Ref. [1] is different from our result [3]. This difference may be due to the choice of parameters in the calculations.

In summary, the damping factor (Eq.(1)) which is commonly applied to analyze the effects of spin, should be used carefully. In the system with magnetic breakdown, some Fourier amplitudes are *enhanced* by spin.

-
- [1] S. Y. Han, J. S. Brooks, and Ju H. Kim, Phys. Rev. Lett. **85**, 1500 (2000).
 - [2] D. Shoenberg, *Magnetic Oscillation in Metals* (Cambridge University Press: Cambridge, 1984).
 - [3] K. Kishigi, Y. Hasegawa and M. Miyazaki, J. Phys. Soc. Jpn. **68** 1817 (1999), **69**, 821 (2000).
 - [4] M. Nakano: Phys. Rev. B **62** 45 (2000).